

Black Sea Ecosystem Modelling

Contents

| | |
|--|---|
| Data provider | 2 |
| Regional summary | 2 |
| Justification of model selection | 3 |
| Technical overview of models used in this region | 4 |
| Hydrodynamic model..... | 4 |
| Lower Trophic level model..... | 4 |
| Higher trophic level models..... | 4 |
| Data Assimilation | 6 |
| Forcing and Boundary Conditions..... | 6 |
| References | 7 |



Data provider


Black Sea models have been developed by the [Institute of Marine Science](#) at the Middle East Technical University (METU) by Dr Baris Salihoglu and his team.

Regional summary

The Black Sea is one of the major semi-enclosed seas of the world oceans and has experienced striking ecological changes due to impacts of climate change, intense eutrophication, population explosion of some invasive species and unsustainable fisheries with their density-dependent feedback processes. Nitrate loading increased nearly four-fold during the 1970s due to increased use of agricultural fertilizers. Rapidly intensifying eutrophication has caused comparable increases both in the subsurface nitrate concentrations and phytoplankton biomass, degradation of the classical mesozooplankton-dominated food web by flourishing of the opportunistic species *Noctiluca scintillans* and *Aurelia aurita* in the 1980s. The intense eutrophication phase was ended by a sequence of events; the collapse of total small pelagic stock (Oguz et al., 2012) and the simultaneous outburst of ctenophore *Mnemiopsis leidyi* population at 1989-1991 (Shiganova et al., 2001; Purcell et al., 2001) and the subsequent marked decline of anthropogenic nutrient loads from the River Danube and other northwestern rivers following the disintegration phase of the former Soviet Union during the early 1990s (Oguz and Velikova, 2010). The subsequent years are referred to as the post-eutrophication phase in which the nutrient reduction led to approximately a two-fold decrease in the phytoplankton and small pelagic standing stocks and *Noctiluca* and *Mnemiopsis* populations (BSC, 2008), and 30-40% decline in the subsurface nitrate and phosphate peak concentrations. Invasion of the opportunistic gelatinous species *Beroe ovata* and their predation on the *Mnemiopsis* population has introduced further changes in the food web structure by the end of 1990s.

Contrary to the relatively rich data availability and their analyses related to the eutrophication and fishery, observations concerning the carbonate system properties are scarce and limited to the total alkalinity and pH measurements. These data sets may be helpful to understand gross features of the carbonate system, but the lack of systematic time series measurements hinders exploring their seasonal to inter-annual variability and the current status of acidification and its vertical structure in the aerobic part of the interior Black Sea.

Different aspects of the structural changes observed in the Black Sea ecosystem have been studied quantitatively by modelling studies (e.g. Oguz et al., 2000, 2001a,b; Oguz and Merico, 2006; Lancelot et al., 2002; Daskalov, 2002; Gregoire and Friedrich, 2004; Gregoire and Soetaert, 2010; Staneva et al., 2010; He et al., 2012). They focussed mostly on the impacts of eutrophication, fishery and invasion of alien gelatinous species, but none of them have investigated the acidification problem in general and the processes governing atmospheric CO₂ uptake, temporal and spatial variability of the carbonate system properties in particular. Therefore, additional modeling studies are timely and warranted to broaden our conceptual understanding on the Black Sea carbonate structure and to analyze the synergistic effects of climate change, eutrophication, and acidification by means of sufficiently complex and fully coupled models of the physical-biogeochemical-carbonate system.




The Black Sea: geographic setting, main rivers and bathymetric features (from Besiktepe et. al., 2001)

Justification of model selection

The peculiar characteristics of the Black Sea that include strong stratification, anoxia below 150m, water and nutrient influx from a high number of major rivers and recent invasion events by alien species make it difficult to model effectively using standard off-the-shelf models. As a result, models that take into account the properties peculiar to the Black Sea were developed at the Institute of Marine Science of the Middle East Technical University (IMS-METU) in previous EU funded projects. The physical, lower-trophic-level and higher-trophic level models were tested and validated extensively during the [MEECE](#) project and their capacity for prediction was duly demonstrated.

Technical overview of models used in this region

Black Sea model is an end-to-end coupled model that includes a physical model (POM), a lower-trophic-level model (BIMS_ECO), a higher-trophic-level box-type model (EwE) and an individual based model larvae model (Anchovy IBM). The physical model is run in a standalone mode to produce the fluid velocity, vertical diffusivity, temperature and salinity fields, which are stored in a NetCDF file that is read by BIMS_ECO to force the biogeochemical dynamics. The biogeochemical fields produced by BIMS_ECO are stored in another NetCDF file that is used in EwE and IBM models as input. All of the couplings in our modelling framework are therefore offline and make use of file I/O for coupling.



The coupling between POM, BIMS_ECO, EwE and Anchovy IBM. The arrows indicate one-way coupling in which the output of one module is stored in a file that is used as input in the other module.

Hydrodynamic model

The hydrodynamic model applied in this study is the Princeton Ocean Model using the Mellor-Yamada level 2.5 turbulence parameterization (POM; Blumberg and Mellor, 1987). The model domain encompasses the entire Black Sea (41°S - 46°N, 28°E - 41.5°E), although excludes the Azov Sea and consists of a (4km x 4km) resolution, Arakawa C, horizontal grid and a 26 level, sigma-coordinate, vertical grid. The sigma-levels are compressed towards the surface and bottom. The maximum depth of the model domain is 220 m.

Lower Trophic level model

The biogeochemical model resolves all the major processes within the strongly stratified upper layer water column that overlies the deep (~2000 m) sulfide-rich anoxic pool. It describes the dynamics of phytoplankton (diatoms, dinoflagellates, small phytoplankton and coccolithophores); bacterioplankton; zooplankton (micro- and mesozooplankton, gelatinous carnivorous zooplankton *Mnemiopsis leidyi* and the heterotrophic dinoflagellate *Noctiluca scintillans*); particulate organic nitrogen and phosphorus; inorganic nutrients (NO₃, NH₄, PO₄); nitrogen gas (N₂); dissolved oxygen and hydrogen sulphide; CaCO₃ produced by coccolithophores; and detritus both in nitrogen and phosphorus currencies. Dissolved CO₂ and total alkalinity (TA) are additional state variables that couple the model to the carbonate module.

Higher trophic level models

Ecosim with Ecopath

The Black Sea HTL model was built to represent the general prevailing food web structure of the Black Sea in the inner basin, prudently avoiding the extremely variable conditions of the Northwestern Shelf (NWS). The model covers an area of 150 000 km² where fisheries operate intensively (Oguz et al., 2008). The geographical representation of the model does not include continental shelf areas and depths greater than 150 m at which anoxia prevails. The mass-balance Ecopath model of the Black Sea was based on Akoglu et al. (in prep.) and further developed by disaggregating the functional groups of small pelagic fish, pelagic piscivorous fish and demersal fish, which were aggregated guilds of multiple species in the earlier study, into individual species. Further, new state variables were introduced; i.e. two new jellyfish species (*Pleurobrachia spp.* and *Beroe ovata*) and another detritus compartment; particulate organic matter (POM). The model was extended to include 23 functional groups, which comprised of 11 fish groups; Black Sea anchovy (*Engraulis encrasicolus ponticus*), Black Sea sprat (*Sprattus sprattus phalaericus*), Pontic shad (*Alosa kessleri pontica*), Black Sea horse mackerel (*Trachurus mediterraneus ponticus*), bonito (*Sarda sarda*), bluefish (*Pomatomus saltator*), Atlantic mackerel (*Scomber scombrus*), turbot (*Psetta maotica*), spiny dogfish (*Squalus acanthias*), Black Sea whiting (*Merlangius merlangus euxinus*) and red mullet (*Mullus barbatus ponticus*), four jellyfish; *Mnemiopsis leidy*, *Aurelia*, *Pleurobrachia spp.* and *Beroe ovate*; two detritus groups; one group representing sediment and one representing particulate organic matter in the water column (POM); one phytoplankton (diatoms and flagellates) and one zooplankton (non-gelatinous, fodder zooplankton) group; one heterotrophic dinoflagellate *Noctiluca scintillans* and one dolphin group to represent the Black Sea marine mammals, which are composed of short-beaked common dolphin (*Delphinus delphis*), bottlenose dolphin (*Tursiops truncatus*) and harbor porpoise (*Phocoena phocoena*). Anchovy was defined as a multi-stanza group separated into “eggs and larvae (0 to 3-month old)” and “juveniles and adults (3 months and above)” stanzas. Three fishing types were identified; trawlers, purse seiners and gill-netters to represent the fishery impact on the ecosystem. The mass-balance model was setup and balanced for the quasi-pristine conditions of the early 1980s. The diet composition matrix of the model was largely based on data available by stomach content analysis and compiled from FishBase .

The initial EwE model of the Black Sea was run 1-way and offline coupled with BIMS-Eco (Oguz et al., 2001a) model. The lower trophic level variables; i.e. phytoplankton, zooplankton, *Aurelia* and *M. leidy* were forced with the simulated biomasses by BIMS-Eco model. The higher trophic level groups; i.e. fish species and dolphins as well as some lower trophic groups; i.e. *Pleurobrachia spp.* which were not included in the BIMS-Eco model, were also simulated by the EwE Black Sea model.

The Ecosim dynamic model was run between 1980 and 2000 for hindcast validation, up until the most recent date at which adequate fisheries biology data were available. The dynamics of the fish compartments were forced using exploitation rates, i.e. fishing mortalities, where available. For groups/species without any fishing mortality estimates and stock assessment values; i.e. red mullet, bluefish and bonito, the catches were forced. Since no catch or stranding statistics were available for marine mammals, i.e. dolphins, in the Black Sea, the fishing mortality for this group was set to an estimated fixed value of 0.5 in order to represent the significant effect of by-catches due to fishery operations. The jellyfish compartments were forced by observed biomass values since they do not have natural predators within the system prior to the introduction of *B. ovata* in 1992. The primary producers were forced with time-series of non-linear production estimates from phytoplankton biomass during the simulation period. The time series of catch data and stock assessments of other models like VPA were used as a measure for the model fitting. The model was calibrated using

vulnerabilities of the functional groups as well as Monte Carlo style parameter searching for the mass-balance Ecopath model.

Data Assimilation

Eight-day composites of Chl-a obtained from the OceanColor product are assimilated into BIMS_ECO every eight days using the Singular Fixed Extended Kalman Filter (SFEK) and the localized Ensemble Transform Kalman Filter (ETKF). Parallel Data Assimilation Framework (PDAF), which is an open-source and freely available software package (<http://pdaf.awi.de/trac/wiki>), is used to implement the aforementioned filters. During the restructuring of BIMS_ECO, the prerequisites for the use of PDAF were considered to ensure compatibility.

Forcing and Boundary Conditions

Hydrodynamic Model

Because the Black Sea is a semi-enclosed basin, the lateral boundary conditions are no-slip and zero heat flux and salt fluxes everywhere except at the mouths of the Bosphorous and Kerch strait, and the 8 largest rivers surrounding the Black Sea (Danube, Dniester, Dnieper, Inguri, Rioni, Yesilirmak, Kizilirmak and Sakarya). At the mouths of the rivers and straits, temperature and salinity boundary conditions are specified as inflow conditions. Diffusive heat and salt fluxes are set to zero in the straits outflow points. Surface forcing is prescribed using 6-hourly fields of wind stress, fresh water fluxes (evaporation, convective precipitation and large-scale precipitation), and radiation fields (surface shortwave radiation, surface long-wave radiation, evaporative heat flux, and convective heat flux) extracted from the DMI HIRHAM regional climate model. All simulations will be augmented by a 5 year spin-up period allowing the model to adjust to the prevalent conditions. Initial conditions are defined using monthly mean climatology for January from the World Ocean Atlas (WOA; Garcia et al., 2006).

Lower Trophic Level Model

Initial conditions. The initial conditions are chosen to reproduce the observed characteristics of the interior Black Sea ecosystem during its post-eutrophication phase. Nitrate concentration is set to 3.0 mmol m⁻³ within the upper 65 m, 9.0 mmol m⁻³ between 65m and 85 m depths and decreasing linearly to zero at 110 m. Phosphate has the same form but its concentrations are an order of magnitude lower. Dissolved oxygen is set to 50 μM within the first 60 m, 1 μM within the next 40 m (including the SOL) and zero further below. All the other state variables of the biogeochemical model are set to small and vertically uniform values over the entire water column because their equilibrium structures do not depend on the initial conditions and are emergent properties of the model dynamics. A spinup with 10 cycles is performed with these initial conditions and using physical and atmospheric forcing for the year 1990.

For full details on models used in this region please refer to http://marine-opec.eu/downloads/OPEC_D2.4.pdf

References

- Akoglu, E., Salihoglu, B., Libralato, S., Oguz, T., Solidoro, C. (in prep.). An indicator-based model assessment of the Black Sea ecosystem structure and functioning during 1960s - 1990s
- Afanasyev, Y.D., A.G. Kostianoy, A.G. Zatsepin, and P.M. Poulain, 2002. Analysis of velocity field in the eastern Black Sea from satellite data during the Black Sea '99 experiment. *J. Geophys. Res.*, 107, 10.1029/2000JC000578.
- Basturk, O., C. Saydam, I. Salihoglu, L. V. Eremeeva, S. K. Konovalov, A. Stoyanov, A. Dimitrov, A. Cociasu, L. Dorogan, and M. Altabet, 1994. Vertical variations in the principle chemical properties of the Black Sea in the autumn of 1991. *J. Marine Chemistry*, 45, 149–165.
- Basturk, O., S.Tugrul, S.Konovalov, I., Salihoglu, 1998a. Effects of circulation on the spatial distributions of principle chemical properties and unexpected short- and long-term changes in the Black Sea. In: *Ecosystem modeling as a management tool for the Black Sea*. Ivanov, L.I. and T. Oguz (eds.), NATO ASI Series, Vol. 47, Kluwer Academic publishers, Netherlands, pp. 39–54.
- Beckers, J.M., M.L. Gregoire, J.C.J. Nihoul, E. Stanev, J. Staneva and C. Lancelot, 2002. Hydrodynamical processes governing exchanges between the Danube, the North-Western continental shelf and the Black Sea's basin simulated by 3D and box models. *Estuarine, Coastal and Shelf Science*, 54, 453–472.
- Besiktepe, S.T., C.J., Lozano, and A.R., Robinson, 2001. On the summer mesoscale variability of the Black Sea. *J. Mar. Res.*, 59, 475–515.
- Beaufort, L., I. Probert, de Garidel-Thoron, T., Bendif, E. M., Ruiz-Pino, D., Metzl, N., Goyet, C., Buchet, N., Coupel, P., Grelaud, M., Rost, B., Rickaby, R. E. M. and de Vargas, C., 2011, Sensitivity of coccolithophores to carbonate chemistry and ocean acidification, *Nature*, 476:80-83.
- Bissett, W.P., Walsh, J.J., Dieterle, D.A. and Carter, K.L., 1999, Carbon cycling in the upper waters of the Sargasso Sea. I. Numerical simulation of differential carbon and nitrogen fluxes, *Deep-Sea Res. I*, 46:205–269.
- BSC (2008). *State of the Environment of the Black Sea (2001-2006/7)*. The Commission on the Protection of the Black Sea Against Pollution publication, 448 pp.
- Codispoti, L.A., G.E. Friederich, J.W. Murray, and C.M. Sakamoto, 1991. Chemical variability in the Black Sea: implications of continuous vertical profiles that penetrated the oxic/anoxic interface. *Deep-Sea Res. I*, 38, Suppl.2, S691-S710.
- Cociasu, A., L. Dorogan, C. Humborg and L. Popa, 1996. Long-term ecological changes in the Romanian coastal waters of the Black Sea. *Marine Pollution Bulletin*, 32, 32–38.
- Codispoti, L.A., G.E. Friederich, J.W. Murray, and C.M. Sakamoto, 1991. Chemical variability in the Black Sea: implications of continuous vertical profiles that penetrated the oxic/anoxic interface. *Deep-Sea Res. I*, 38, Suppl.2, S691-S710.
- Daskalov GM. Overfishing drives a trophic cascade in the Black Sea. *Mar. Ecol. Progr. Ser.*, 2002; 225: 53–63.
- Di Iorio, D. and H. Yuce, 1998. Observations of Mediterranean flow into the Black Sea. *J. Geophys. Res.*, 104, 3091–3108.
- Dyrssen, D.W. (1999). Framvaren and the Black Sea – Similarities and Differences. *Aquatic Geochemistry* 5: 59–73.
- Egge, J.K, Thingstad, T.F., Larsen, A., Engel, A., Wohlers, J., Bellerby, R.G.J. and Riebesell, U. 2009,

Primary production during nutrient-induced blooms at elevated CO₂ concentrations, *Biogeosciences*, 6:877-885.

Engel, A., Schulz, K. G., Riebesell, U., Bellerby, R., Delille, B. and Schartau, M., 2008, Effects of CO₂ on particle size distribution and phytoplankton abundance during a mesocosm bloom experiment (PeECE II), *Biogeosciences*, 5:509–521.

Fasham, M. J. R., Ducklow, H. W. and McKelvie, S.M., 1990, A nitrogen-based model of plankton dynamics in the oceanic mixed layer, *J. Mar. Res.*, 48:591–639.

Feng, Y., Warner, M. E., Zhang, Y., Sun, J., Fu, F., Rose, J. M. and Hutchins, D. A., 2008, Interactive effects of increased pCO₂, temperature and irradiance on the marine coccolithophore *Emiliana huxleyi* (Prymnesiophyceae), *Eur. J. Phycol.*, 43:87–98.

Gangstø, R., Gehlen, M., Schneider, B., Bopp, L., Aumont, O. and Joos, F., 2008, Modeling the marine aragonite cycle: changes under rising carbon dioxide and its role in shallow water CaCO₃ dissolution, *Biogeosciences*, 5:1057–1072.

Garcia A, Palomera I (1996) Anchovy early life history and its relation to its surrounding environment in the Western Mediterranean basin. *Sci. Mar.* 60 (Supl. 2): 155-166.

Gawarkiewicz, G., G. Korotaev, S. Stanichny, L. Repetin, and D. Soloviev, 1999. Synoptic upwelling and cross-shelf transport processes along the Crimean coast of the Black Sea. *Cont. Shelf Res.*, 19, 977–1005.

Gehlen, M., Gangstø, R., Schneider, B., Bopp, L., Aumont, O. and Ethe, C., 2007, The fate of pelagic CaCO₃ production in a high CO₂ ocean: A model study, *Biogeosciences*, 4:505–519.

Gentleman, W., Leising, A., Frost, B., Strom, S. and Murray, J., 2003, Functional responses for zooplankton feeding on multiple resources: a review of assumptions and biological dynamics, *Deep-Sea Res. II*, 50:2847–2875.

Ginsburg, A.I., A.G., Kostianoy, N.P., Nezlin, D.M., Soloviev, S.V., Stanichny, 2002a. Anticyclonic eddies in the northwestern Black Sea. *J. Mar. Syst.*, 32, 91–106.

Ginsburg, A.I., A.G., Kostianoy, V.G., Krivosheya, N.P., Nezlin, D.M., Soloviev, S.V., Stanichny, V.G. Yakubenko, 2002b. Mesoscale eddies and related processes in the northeastern Black Sea. *J. Mar. Syst.*, 32, 71–90.

Goyet, C., A. L. Bradshaw, and P. G. Brewer (1991), The carbonate system in the Black Sea, *Deep-Sea Res.*, 38, No. 2, 1049–1067.

Gregg, M.C. and E. Ozsoy, 1999. Mixing on the Black Sea shelf north of the Bosphorus. *Geophys. Res. Letters*, , 26, 1869–1872.

Grégoire, M., Friedrich, J., 2004. Nitrogen budget of the north-western black sea shelf as inferred from modeling studies and in-situ benthic measurements. *Marine Ecology Progress Series* 270, 15–39.

Grégoire, M. and K. Soetaert (2010) Carbon, nitrogen, oxygen and sulfide budgets in the Black Sea: A biogeochemical model of the whole water column coupling the oxic and anoxic parts. *Ecological Modelling*, 221, 2287–2301.

He, Y., E. V. Stanev, E. Yakushev, J. Staneva (2012) Black Sea biogeochemistry: Response to decadal atmospheric variability during 1960-2000 inferred from numerical modeling. *Marine Environmental Research*, 77, 90-102.

Hiscock, W.T., and F. J. Millero (2006) Alkalinity of anoxic waters in the Western Black Sea. *Deep-Sea*

Res., 53, 1787–1801.

Ilyina, T., Zeebe, R. E., Maier-Reimer, E. and Heinze, C., 2009, Early detection of ocean acidification effects on marine calcification, *Global Biogeochemical Cycles*, 23, GB1008, doi:10.1029/2008GB003278.

Ivanov, L.I. and A.S. Samodurov, 2001. The role of lateral fluxes in ventilation of the Black Sea. *J. Mar. Syst.*, 31, 159–174.

Joassin, P., Delille, B., Soetaert, K., Harlay, J., Borges, A. V., Chou, L., Riebesell, U., Suykens, K. and Grégoire, M., 201, Carbon and nitrogen flows during a bloom of the coccolithophore *Emiliania huxleyi*: Modelling a mesocosm experiment, *J. Mar. Syst.*, 85:71–85.

Karl, D.M., and G.A. Knauer, 1991. Microbial production and particle flux in the upper 350 m of the Black Sea. *Deep Sea Res.*, 38, Supp. 2A, S655-S661.

Kideys A.E. (2002) Fall and rise of the Black Sea ecosystem. *Science* 297: 1482-1484.

Kideys AE, Gordina AD, Bingel F, Niermann U (1999) The effect of environmental conditions on the distribution of eggs and larvae of anchovy (*Engraulis encrasicolus* L.) in the Black Sea. *ICES J. Mar. Sci.* 56 (Supplement): 58-64.

Konovalov, S.K., and J.W. Murray, 2001. Variations in the chemistry of the Black Sea on a time scale of decades (1960–1995). *J. Mar. Syst.*, 31, 217–243.

Konovalov, S.K., G.W. Luther, III, G.E. Friederich, D.B. Nuzzio, B.M. Tebo, J.W. Murray, T. Oguz, B. Glazer, R.E. Trouwborst, B. Clement, K. W. Murray, A. S. Romanov, 2003. Lateral injection of oxygen with the Bosphorus plume-fingers of oxidizing potential in the Black Sea. *Limnol. Oceanogr.*, 48, 2369–2376.

Korotaev, G.K., T. Oguz, A. Nikiforov, C. J. Koblinsky, 2003. Seasonal, interannual and mesoscale variability of the Black Sea upper layer circulation derived from altimeter data. *J. Geophys. Res.*, 108(C4), 3122, doi:10.1029/2002JC001508.

Krivosheya, V.G., V. B. Titov, I.M. Ovchinnikov, R.D. Kosyan, A.Yu. Skirta, 2000. The influence of circulation and eddies on the depth of the upper boundary of the hydrogen sulfide zone and ventilation of aerobic waters in the Black Sea. *Oceanology (Eng. Transl.)*, 40, 767–776.

Krivosheya, V.G., I.M. Ovchinnikov, and A. Yu. Skirta, 2002. Interannual variability of the cold intermediate layer renewal in the Black Sea. In: *Multidisciplinary investigations of the northeast part of the Black Sea*, A.G. Zatsepin and M.V. Flint, eds, Moscow, Nauka, pp. 27–39.

Kuypers, M.M.M., A.O. Sliemers, G. Lavik, M. Schmid, B.B. Jorgensen, J.G. Kuenen, D.J.S. Sinninghe, M. Strous, and M.S.M. Jetten, 2003. Anaerobic ammonium oxidation by anammox bacteria in the Black Sea. *Nature*, 422, 608–611.

Lancelot C, Staneva J, Van Eeckhout D, Beckers JM, Stanev E (2002) Modeling the impact of the human forcing on the ecological functioning of the northwestern Black Sea. *Estuarine, Coastal and Shelf Science* 54: 473-500.

Latif, M.A., E. Ozsoy, T. Oguz, and U. Unluata, 1991. Observation of the Mediterranean inflow into the Black Sea. *Deep-Sea Research*, 38, Supl.2A, S711-S733.

Lisovenko LA, Andrianov DP (1996) Reproductive biology of anchovy (*Engraulis encrasicolus ponticus* Alexandrov 1927) in the Black Sea. *Sci. Mar.* 60 (suppl): 209-218.

Lebedeva, L.P. and S. V. Vostokov, 1984. Studies of detritus formation processes in the Black Sea. *Oceanology (Engl. Transl.)*, 24(2), 258–263.

- Luo J, Brandt SB (1993) Bay anchovy *Anchoa mitchilli* production and consumption in mid-Chesapeake Bay based on a bioenergetic model and acoustic measures of fish abundance. *Mar. Ecol. Prog. Ser.* 98: 223-236.
- Megrey BA, Rose KA, Klumb RA, Hay DE, Werner FE, Eslinger DL, Smith SL (2007) A bioenergetic-based population dynamics model of pacific herring (*Clupea harengus pallasi*) coupled to a lower trophic level nutrient-plankton-zooplankton model: Description, calibration, and sensitivity analysis. *Ecol. Model.* 202: 144-164.
- Mehrbach, C., Culberson, C. H., Hawley, J. E. and Pytkowicz, R.M., 1973, Measurement of the apparent dissociation constants of carbonic acid in seawater at atmospheric pressure, *Limnol. Oceanogr.*, 18:897–907.
- Merico, A., Tyrrell, T. and Cokacar, T., 2006, Is there any relationship between phytoplankton seasonal dynamics and the carbonate system?, *J. Mar. Syst.*, 59:120–142.
- Moiseenko, O. G., S. K. Konovalov, and O. N. Kozlovskaya (2011) Intra-annual and long-term variations of the carbonate system of the aerobic zone in the Black Sea. *Physical Oceanography*, 20, 435-449.
- Mucci, A., 1983, The solubility of calcite and aragonite in seawater at various salinities, temperatures, and one atmosphere total pressure, *Am. J. Sci.*, 283:781–799.
- Murray, J.W., H. W. Jannash, S. Honjo, R. F. Anderson, W.S. Reeburgh, Z. Top, G.E. Friederich, L.A. Codispoti, and E. Izdar, 1989. Unexpected changes in the oxic/anoxic interface in the Black Sea. *Nature*, 338, 411–413.
- Murray, J. W., Z. Top, and E. Ozsoy, 1991. Hydrographic properties and ventilation of the Black Sea. *Deep-Sea Res.*, 38, Suppl.2A, S663–690.
- Murray, J.W., H. W. Jannash, S. Honjo, R. F. Anderson, W.S. Reeburgh, Z. Top, G.E. Friederich, L.A. Codispoti, and E. Izdar, 1989. Unexpected changes in the oxic/anoxic interface in the Black Sea. *Nature*, 338, 411–413.
- Murray, J.W., L.A. Codispoti, and G.E. Friederich, 1995. Oxidation-reduction environments: The suboxic zone in the Black Sea. In: *Aquatic chemistry: Interfacial and interspecies processes*. C.P. Huang, C.R.O’Melia, and J.J. Morgan, eds. ACS Advances in Chemistry Series No.224. pp. 157–176.
- Murray, J.W., B-S Lee, J. Bullister, and G. W. Luther III, 1999. The suboxic zone of the Black Sea. In: *Environmental Degradation of the Black Sea: Challenges and Remedies*. S. Besiktepe, U. Unluata and A. Bologna, eds. NATO ASI Series 2., 75–92.
- Nezlin, N.P., 2001. Unusual phytoplankton bloom in the Black Sea during 1998–1999: Analysis of remotely-sensed data, *Oceanology*, (English transl.), 41, 375–380.
- Niermann U, Bingel F, Gorban A, Gordina AD, Gücü AC, Kideys AE, Konsulova A, Radu G, Subbotin AA, Zaika VE (1994) Distribution of anchovy eggs and larvae (*Engraulis encrasicolus* Cuv.) in the Black Sea in 1991-1992. *ICES J. Mar. Sci.* 51:395-406.
- Neumann, T, 2000, Towards a 3D-ecosystem model of the Baltic Sea, *J. Mar. Syst.*, 25:405-419.
- North EW, Houde ED (2004) Distribution and transport of bay anchovy (*Anchoa mitchilli*) eggs and larvae in Chesapeake Bay. *Estuarine, Coastal and Shelf Science* 60:409-429.
- Oguz, T. and L. Rozman, 1991. Characteristics of the Mediterranean underflow in the southwestern Black Sea continental shelf/slope region. *Oceanology Acta*, 14(5), 433-444.
- Oguz, T., S., Besiktepe, 1999. Observations on the Rim Current structure, CIW formation and

transport in the western Black Sea. *Deep Sea Res.*, 46, 1733-1753.

Oguz, T. and Ediger, D., 2006. Comparison of in-situ and satellite-derived chlorophyll pigment concentrations and impact of phytoplankton bloom on the suboxic layer structure in the western Black Sea during May-June 2001. *Deep Sea Research Part II: Topical Studies in Oceanography*, 53 (17-19): 1923-1933.

Oguz, T., E. Ozsoy, M.A. Latif, H.I. Sur and U. Unluata, 1990a. Modeling of hydrographically controlled exchange flow in the Bosphorus Strait. *J. Phys. Oceanogr.*, 20, 945–965.

Oguz, T., M.A. Latif, H. I. Sur, E. Ozsoy, and U. Unluata, 1990b. On the dynamics of the southern Black Sea. In: *The Black Sea Oceanography*, J. Murry and E. Izdar, eds, NATO/ASI Series, Kluwer Academic Publishers, Dordrecht, pp. 43–64.

Oguz, T., P., La Violette, U., Unluata, 1992. Upper layer circulation of the southern Black Sea: Its variability as inferred from hydrographic and satellite observations. *J. Geophys. Res.*, 97, 12569–12584.

Oguz, T., V.S., Latun, M.A., Latif, V.V., Vladimirov, H.I., Sur, A.A., Makarov, E., Ozsoy, B.B., Kotovshchikov, V.V., Ereemeev, U., Unluata, 1993. Circulation in the surface and intermediate layers of the Black Sea. *Deep Sea Res. I*, 40, 1597–1612.

Oguz, T., D.G., Aubrey, V.S., Latun, E., Demirov, L., Koveshnikov, H.I., Sur, V., Diacanu, S., Besik-tepe, M., Duman, R., Limeburner, V.V., Ereemeev, 1994. Mesoscale circulation and thermohaline structure of the Black sea observed during HydroBlack'91. *Deep Sea Research I*, 41, 603–628.

Oguz, T., P., Malanotte-Rizzoli, D., Aubrey, 1995. Wind and thermohaline circulation of the Black Sea driven by yearly mean climatological forcing. *J. Geophys. Res.*, 100, 6846–6865.

Oguz, T., S. Tugrul, A.E. Kideys, V. Ediger, N. Kubilay (2005) "Physical and biogeochemical characteristics of the Black Sea". *The Sea*, Vol. 14, Chapter 33, 1331-1369.

Oguz, T., T. Cokacar, P. Malanotte-Rizzoli, and H. W. Ducklow, 2003. Climatic warming and accompanying changes in the ecological regime of the Black Sea during 1990s. *Global Biogeochem. Cycles*, 17(3), 1088, doi:10.1029/2003.

Oguz, T., L.I. Ivanov, S. Besiktepe, 1998. Circulation and hydrographic characteristics of the Black Sea during July 1992. In: *Ecosystem Modeling as a Management Tool for the Black Sea*, L. Ivanov and T. Oguz (eds). NATO ASI Series, Environmental Security-Vol.47, Kluwer Academic Publishers, Vol. 2, pp. 69–92.

Oguz, T., H. W. Ducklow, and P. Malanotte-Rizzoli, 2000. Modeling distinct vertical biogeochemical structure of the Black Sea: Dynamical coupling of the Oxidic, Suboxic and Anoxic layers. *Global Biogeochem. Cycles.*, 14, 1331–1352.

Oguz, T., A. G. Deshpande, and P. Malanotte-Rizzoli, 2002a. On the Role of Mesoscale Processes Controlling Biological Variability in the Black Sea: Inferences From SeaWiFS-derived Surface Chlorophyll Field. *Continental Shelf Research*, 22, 1477–1492.

Oguz, T., and A. Merico (2006) Factors controlling the summer *Emiliania huxleyi* bloom in the Black Sea: a modeling study. *J. Marine Systems*, 59, 173-188.

Oguz T and Velikova V. 2010, Abrupt transition of the northwestern Black Sea shelf ecosystem from a eutrophic to an alternative pristine state. *Mar. Ecol. Prog. Ser.*, 405: 231–242.

Oguz T, Salihoglu B, Moncheva S, Abaza V. 2012b, Regional peculiarities of community-wide trophic cascades in strongly degraded Black Sea food web. *J. Plankton Res.*, 34: 338-343.

- Oguz, T., Fach, B., Salihoglu, B., 2008. A coupled plankton-anchovy population dynamics model assessing nonlinear controls of anchovy and gelatinous biomass in the Black Sea. *Mar Ecol Prog Ser*, Vol. 369, 229-256.
- Oguz, T., J. W. Murray, and A. E. Callahan, 2001a. Modeling redox cycling across the suboxic-anoxic interface zone in the Black Sea. *Deep-Sea Res. I.*, 48, 761–787.
- Oguz, T., P. Malanotte-Rizzoli, H.W. Ducklow (2001b) Simulations of phytoplankton seasonal cycle with multi-level and multi-layer physical-ecosystem models: The Black Sea example. *Ecological Modelling*, 144, 295-314.
- Oguz, T. , Arkin., S. S., Akoglu, E., Fach, B., Salihoglu, B. (2013) Modeling intrinsic controls of the carbonate system and enlargement of acidification by eutrophication in the Black Sea (Submitted to MEPS)
- O'Neill, R. V., Angelis, D. L., Pastor, J. J., Jackson, B. J. and Post, W. M., 1989, Multiple nutrient limitation in ecological modeling, *Ecol. Model.*, 46:147–163.
- Ozsoy, E., U. Unluata and Z. Top, 1993. The Mediterranean water evolution, material transport by double diffusive intrusions and interior mixing in the Black Sea. *Prog. Oceanogr.*, 31, 275–320.
- Ozsoy, E., U. Unluata and Z. Top, 1993. The Mediterranean water evolution, material transport by double diffusive intrusions and interior mixing in the Black Sea. *Prog. Oceanogr.*, 31, 275–320.
- Ozsoy, E., D. Di Iorio, M. C. Gregg, and J.O. Backhaus, 2001. Mixing in the Bosphorus Strait and the Black Sea continental shelf: observations and a model of dense water outflow. *J. Mar. Syst.*, 31, 99–135.
- Ozsoy, E., D. Rank, and I. Salihoglu, 2002. Pycnocline and deep mixing in the Black Sea: stable isotope and transient tracer measurements. *Estuarine, Coastal and Shelf Science*, 54, 621–629.
- Paulmier, A., Kriest, I. and Oeschies, A., 2009, Stoichiometries of remineralisation and denitrification in global biogeochemical ocean models, *Biogeosciences*, 6:923–935.
- Peneva, E., E. Stanev, V. Belokopytov, and P.Y. Le Traon, 2001. Water transport in the Bosphorus Strait estimated from hydro-meteorological and altimeter data: seasonal and decadal variability. *J. Mar. Syst.*, 31, 21–33.
- Polat, S.Ç. and S., Tugrul, 1995. Nutrient and organic carbon exchanges between the Black and Marmara Seas through the Bosphorus Strait. *Continental Shelf Research*, 15, 1115–1132.
- Purcell, J.E., Shiganova, T.A., Decker, M.B. and Houde, E.D. (2001) The ctenophore *Mnemiopsis* in native and exotic habitats: U.S. estuaries versus the Black Sea basin. *Hydrobiologia*, 451, 145-176.
- Repeta, D.J. and D.J. Simpson, 1991. The distribution and recycling of chlorophyll, bacteriochlorophyll and carotenoids in the Black Sea. *Deep-Sea Res.*, 38, Suppl.2A, S969-S984.
- Riebesell, U., et al., 2007, Enhanced biological carbon consumption in a high CO₂ Ocean, *Nature*, 450:545–548.
- Repeta, D.J., D.J. Simpson, B.B Jorgensen, and H.W. Jannash, 1989. Evidence for anoxic photosynthesis from distribution of bacteriochlorophylls in the Black Sea. *Nature*, 342, 69–72.
- Romaniello, S. J. and Derry, L. A., 2010, Validation of an intermediate complexity model for simulating marine biogeochemistry under anoxic conditions in the modern Black Sea, *Geochem. Geophys. Geosyst* Volume 11, 8, Q08002, doi:10.1029/2009GC002712.
- Rose KA, Cowan JH, Clark ME, Houde ED, Wang SB (1999) An individual-based model of bay anchovy

population dynamics in the mesohaline region of Chesapeake Bay. *Mar. Ecol. Prog. Ser.* 185: 113-132.

Rozanov, A.G. 1996. Redox stratification in Black Sea waters. *Oceanology (Engl. Transl.)*, 35, 500–504.

Satilmis HH, Gordina AD, Bat L, Bircan R, Culha M, Akbulut M, Kideys AE (2003) seasonal distribution of fish eggs and larvae off Sinop (the southern Black Sea) in 1999-2000. *Acta Oecologica* 24: S275-S280.

Saydam, C., S. Tugrul, O. Basturk, and T. Oguz, 1993: Identification of the oxic/anoxic interface by isopycnal surfaces in the Black Sea. *Deep Sea Research*, 40, 1405–1412.

Shaffer, G., 1986: Phosphate pumps and shuttles in the Black Sea. *Nature*, 321, 515–517.

Shiganova TA, Mirzoyan ZA, Studenikina EA, et al. (2001) Population development of the invader ctenophore *Mnemiopsis leidyi* in the Black Sea and other seas of the Mediterranean basin. *Mar. Biol.*, 139: 431–445.

Soetaert, K., Hofmann, A. F., Middelburg J. J., Meysman F. J. R. and Greenwood, J., 2007, The effect of biogeochemical processes on pH, *Mar. Chem.*, 105:30–51.

Sokolova, E., E.V. Stanev, V. Yakubenko, I. Ovchinnikov, R. Kosyan, 2001. Synoptic variability in the Black Sea. Analysis of hydrographic survey and altimeter data. *J. Mar. Syst.*, 31, 45–63.

Stanev, E. V. and N.H. Rachev, 1999. Numerical study of the planetary Rossby modes in the Black Sea. *J. Mar. Syst.*, 21, 283–306.

Stanev, E.V. and E. Peneva, 2002. Regional sea level response to global climatic change: Black Sea examples. *Global and Planetary Changes*, 32, 33-47.

Staneva, J.V. and E.V. Stanev, 2002. Water mass formation in the Black Sea during 1991–1995. *J. Mar. Syst.*, 32, 199–218.

Staneva, J.V. and E.V. Stanev, 2002. Water mass formation in the Black Sea during 1991–1995. *J. Mar. Syst.*, 32, 199–218.

Staneva, J., V. Kourafalou, and K. Tsiaras (2010) Seasonal and Interannual Variability of the North-Western Black Sea Ecosystem. *Terr. Atmos. Ocean. Sci.*, 21, 163-180.

Sur, H.I., E., Ozsoy, U., Unluata, 1994. Boundary current instabilities, upwelling, shelf mixing and eutrophication processes in the Black Sea. *Progr. Oceanogr.*, 33, 249–302.

Sur, H.I., E., Ozsoy, Y.P., Ilyin, U., Unluata, 1996. Coastal/deep ocean interactions in the Black Sea and their ecological/environmental impacts. *J. Marine Systems*, 7, 293–320, 1996.

Sur, H.I. and Y.P. Ilyin, 1997. Evolution of satellite derived mesoscale thermal patterns in the Black Sea. *Prog. Oceanogr.*, 39, 109–151.

Sur, H.I., E., Ozsoy, U., Unluata, 1994. Boundary current instabilities, upwelling, shelf mixing and eutrophication processes in the Black Sea. *Progr. Oceanogr.*, 33, 249–302.

Tarifeno E, Carmona M, Llanos-Rivera A, Castro LR (2008) Temperature effects on the anchoveta *Engraulis ringens* egg development: do latitudinal differences occur? *Environ. Biol. Fish.* 81:387-395.

Titov, V.B., 2000. Dependence of the formation of the winter hydrological structure in the Black Sea on the severity of winter conditions. *Oceanology (Engl. Transl.)*, 40, 777–783.

Tugrul, S., O. Basturk, C. Saydam, and A. Yilmaz, 1992. The use of water density values as a label of

chemical depth in the Black Sea. *Nature*, 359, 137–139.

UNESCO (1996) Progress on oceanographic tables and standards 1983-1986: work and recommendations of the UNESCO/SCOR/ICES/IAPSO Joint Panel, *Tech. Pap. Mar. Sci.*, 50, p. 50.

Unluata, U., T. Oguz, M.A. Latif and E. Ozsoy, 1989. On the physical oceanography of Turkish straits. In: *The physical oceanography of sea straits*, L.J. Pratt, eds. NATO ASI Series, Kluwer, The Netherlands.

Vasas, V., Lancelot, C., Rousseau, V. and Jordán, F., 2007, Eutrophication and overfishing in temperate nearshore pelagic food webs: a network perspective., *Mar. Ecol. Prog. Ser.*, 336:1–14.

Wanninkhof, R., 1992, Relationship between wind-speed and gas-exchange over the ocean. *Journal of Geophysical Research-Oceans*, 97(C5):7373–7382.

Weiss, R. F., 1974, Carbon dioxide in water and seawater: the solubility of a non-ideal gas, *Mar. Chem.*, 2:203–215.

Yakushev, E.V., D.E. Besedin, Yu. F. Lukashev, and V.K. Chasovnikov, 2001. On the rise of the upper boundary of the anoxic zone in the density field of the Black Sea. *Oceanology (Eng. Transl.)*, 41, 654–659.

Yakushev, E., Pollehne, F., Jost, G., Kuznetsov, I., Schneider, B., Umlauf, L., 2007. Analysis of the water column oxic/anoxic interface in the black and baltic seas with a numerical model. *Marine Chemistry* 107, 388–410.

Yilmaz, A., S. Tugrul, C. Polat, D. Ediger, Y. Coban, and E. Morkoc, 1998a. On the production, elemental composition (C,N,P) and distribution of photosynthetic organic matter in the Southern Black Sea. *Hydrobiologia*, 363, 141–156.